

CONSTRAINT RELAXATION

PEDRO MARRONETTI

*Florida Atlantic University,
777 Glades Rd, Boca Raton, FL 33431, USA
pmarrone@fau.edu*

Full relativistic simulations in three dimensions are known to develop runaway modes that grow exponentially and are accompanied by violations of the Hamiltonian and momentum constraints. We present here a method that controls the violation of these constraints and is tested with simulations of binary neutron stars in circular orbits. We show that this technique improves the overall quality of the simulations.



1. Introduction

For most of the past decade, the main theoretical thrust in gravitational research has been directed toward obtaining stable and accurate numerical models of compact-object binary systems. One of the most difficult problems to tackle has been the control of exponentially growing instabilities that degrade the quality of any simulation and, eventually, terminate it. We propose in Refs. ^{1,2} an evolution scheme (Constraint relaxation or CR) where the Hamiltonian and momentum constraints are only approximately solved at every time step, gently steering the evolution toward the space of solutions of the Einstein Field Equations without completely forcing their numerical satisfaction. This method utilizes the conformal decomposition of the spatial metric and extrinsic curvature, which has traditionally been used to solve the initial value problem for binary systems. In this decomposition, a conformal factor ψ factored out of the spatial metric and a longitudinal addition to the extrinsic curvature generated from a vector potential w_i are used to satisfy the Hamiltonian and the three components of the momentum constraint respectively. CR drives ψ to the solution space of the Hamiltonian constraint by means of a parabolic equation for the conformal factor. Similarly, the momentum constraint is controlled by the use of w_i to push the simulation toward the space of solutions of the momentum constraint. In both cases, a full relaxation of ψ and w_i would lead to the numerical solution of the constraints. However, the stability of the relaxation methods relies on gently updating these fields during the evolution. We showed in Ref.² that a full relaxation scheme becomes unstable rather quickly when used in combination with BSSN.

2. Results

We tested the CR scheme with simulations of an irrotational BNS system. The details of the initial data set are provided in table 1 of Ref. ¹ and the corresponding convergence tests are presented in Ref. ² The plots show curves that, for clarity, have been normalized to their corresponding initial values.

The evolution of the L_2 norm of the Hamiltonian constraint residual $\mathcal{H}\mathcal{C}$ and the three components of the momentum constraint residual $\mathcal{M}_i$ across the numerical

Fig. 1. Evolution of the L_2 norm of the Hamiltonian constraint violation.Fig. 2. Evolution of the L_2 norm of the momentum constraint violation.

grid are shown in Figs. 1 and 2 respectively. CR achieves an impressive reduction of the Hamiltonian constraint violation (Fig.1). Note that the curves are plotted in logarithmic scale to highlight the more than two orders of magnitude difference between the CR (dashed) and BSSN (solid) results. CR not only suppresses the constraint violation modes, but also reduces the violation present in the ID set by

a factor of about 5. The difference between the CR (solid) and the BSSN (dashed) results in the case of the momentum constraint (Fig. 2) is not that impressive. At the end of the simulation, the momentum constraint violation was about four times smaller than in the BSSN runs. The spikes present in the BSSN curves at $t \simeq 0.4P$ occur on the stellar surface and are related to matter displacement in the grid, a side-effect of using a frozen shift vector. Note, however, that those spikes disappear when using CR.




Figure 3 shows the evolution of the total angular momentum. The plot shows the CR (solid), BSSN (dashed) curves, as well as the PN estimation (dotted line) of the angular momentum loss for a point-mass binary with the same mass and angular momentum as the BNS in consideration (see Appendix B of Ref. ¹). The CR based run agrees with the PN prediction for about 1.5 orbital periods. The inset of figure 3 zooms in on the first half of the period, showing the reduced level of noise in the CR curve.

Acknowledgments

This work was partially supported by the National Science Foundation under grant PHY-0555644 and the National Computational Science Alliance under grants PHY020007N, PHY050010T, and PHY050015N.

References

1. P. Marronetti, *Class. Quant. Grav.* **22**, 2433 (2005). gr-qc/0501043.
2. P. Marronetti, *Class. Quant. Grav.* **23**, 2681 (2006). gr-qc/0511143.